Microbiological Safety of Fresh and Fresh-Cut Melons: A Bibliography

Complied by

Robert L. Buchanan, Ph.D. Center for Food Safety and Security Systems College of Agriculture and Natural Resources University of Maryland 0119 Symons Hall College Park, Maryland, USA

July 1, 2016

Gayler, G.E. et al. An outbreak of salmonellosis traced to watermelon. Public Health Record 70:311-313. 1955.

Larson, A. et al. *Salmonella oranienberg* gastroenteritis associated with consumption of pre-cut watermelon – Illinois. Morbidity Mortality Weekly Report 28:522-523. 1979.

Blostein, J. An outbreak of *Salmonella* Javiana associated with consumption of watermelon. Journal of Environmental Health 56(1):29-31. 1993.

Golden, D.A. et al. Growth of *Salmonella* spp. in cantaloupe, watermelon, and honeydew melons. Journal of Food Protection 56:194-196. 1993.

Castillo, A. and Escartin, E.F. Survival of *Campylobacter jejuni* on sliced watermelon and papaya. Journal of Food Protection 57:166-168. 1994.

Herwaldt, B.L. et al. Characterization of a variant strain of Norwalk virus from a food-borne outbreak of gastroenteritis on a cruise ship in Hawaii. Journal of Clinical Microbiology 32:861-866. 1994.

Del Rosariao, B.A. and Beuchat, L.R. Survival and growth of enterohemorrhagic *Escherichia coli* O157:H7 in cantaloupe and watermelon. Journal of Food Protection 58:105-107. 1995.

Beuchat, L.R. Pathogenic microorganisms associated with fresh produce. Journal of Food Protection 59:204-216. 1996.

Bradford, M.A. et al. The cross-contamination and survival of *Salmonella enteritidis* PT4 on sterile and non-sterile foodstuffs. Letters in Applied Microbiology 24:261-264. 1997.

Ayhan, Z. and Chism, G.W. The shelf-life of minimally processed fresh cut melons. Journal of Food Quality 21:29-40. 1998.

Brackett, R.E. Incidence, contributing factors, and control of bacterial pathogens in produce. Postharvest Biology and Technology 15:305-311. 1999.

Larson, A.E. and Johnson, E.A. Evaluation of botulinal toxin production in packaged fresh-cut cantaloupe and honeydew melons. Journal of Food Protection 62:948-952. 1999.

Mohle-Boetani, J.C. et al. An outbreak of *Salmonella* serogroup Saphra due to cantaloupes from Mexico. Journal of Infectious Diseases. 180:1361-1364. 1999.

Park, C.M. and Beuchat, L.R. Evaluation of sanitizers for killing *Escherichia coli* O157:H7, *Salmonella*, and naturally occurring microorganisms on cantaloupes, honeydew melons, and asparagus. Dairy, Food and Environmental Sanitation 19:842-847. 1999.

Leverentz, B. et al. Examination of bacteriophage as a biocontrol method for *Salmonella* on fresh-cut fruit: A model study. Journal of Food Protection 64:1116-1121. 2001.

Sapers, G.M. et al. Antimicrobial treatments for minimally processed cantaloupe melon. Journal of Food Science 66:345-349. 2001.

Shearer, A.E.H. et al. Evaluation of a polymerase chain reaction-based system for detection of *Salmonella* Enteritidis, *Escherichia coli* O157:H7, *Listeria* spp., and *Listeria monocytogenes* on fresh fruits and vegetables. Journal of Food Protection 64:788-795. 2001.

Ukuku, D.O. and Sapers, G.M. Effect of sanitizer treatments on *Salmonella* Stanley attached to the surface of cantaloupe and cell transfer to fresh-cut tissues during cutting practices. Journal of Food Protection 64:1286-1291. 2001.

Ukuku, D.O. et al. Influence of washing treatment on native microflora and *Escherichia coli* population of inoculated cantaloupes. Journal of Food Safety 21:31-47. 2001.

Ukuku, D.O. et al. Bioluminescence ATP assay for estimating total plate counts of surface microflora of whole cantaloupe and determining efficacy of washing treatments. Journal of Food Protection 64:813-819. 2001.

Anderson, S.M. et al. Multistate outbreaks of *Salmonella* serotype Poona infections associated with eating cantaloupe from Mexico --- United States and Canada, 2000—2002. Morbidity and Mortality Weekly Report 51:1044-1047. 2002.

Kozempel, M. et al. Application of the vacuum/steam/vacuum surface intervention process to reduce bacteria on the surface of fruits and vegetables. Innovative Food Science and Emerging Technologies 3:62-72. 2002.

Roller, S. and Seedhar, P. Carvacol and cinnamic acid inhibit microbial growth in freshcut melon and kiwifruit at 4° and 8°C. Letters in Applied Microbiology 35:390-394. 2002.

Ukuku, D.O. and Fett, W.F. Effectiveness of chlorine and nisin-EDTA treatments of whole melons and fresh-cut pieces for reducing native microflora and extending shelf-life. Journal of Food Safety 22:231-253. 2002.

Ukuku, D.O. and Fett, W.F. Behavior of *Listeria monocytogenes* inoculated on cantaloupe surfaces and efficacy of washing treatments to reduce transfer from rind to fresh-cut pieces. Journal of Food Protection 65:924-930. 2002.

Ukuku, D.O. and W.F. Fett. Relationship of cell surface charge and hydrophobicity to strength of attachment of bacteria to cantaloupe rind. Journal of Food Protection 65:1093-1099. 2002.

Aguayo, E. et al. Keeping quality and safety of minimally fresh processed melon. European Food Research and Technology 216:494-499. 2003.

Barak, J.D. et al. Recovery of surface bacteria from and surface sanitization of cantaloupes. Journal of Food Protection 66:1805-1810. 2003.

Behrsing, J. Survival of *Listeria innocua*, *Salmonella salford* and *Escherichia coli* on the surface of fruit with inedible skins. Postharvest Biology and Technology 29:249-256. 2003.

Caldwell, K.N. et al. Attraction of a free-living nematode, *Caenorhabditis elegans*, to foodborne pathogenic bacteria and its potential as a vector of *Salmonella* Poona for preharvest contamination of cantaloupe. Journal of Food Protection 66:1964-1971. 2003.

Gagliardi, J.V. et al. On-farm and postharvest processing of bacterial contamination to melon rinds. Journal of Food Protection 66:82-87. 2003.

Leverentz, B. et al. Biocontrol of *Listeria monocytogenes* on fresh-cut produce by treatment with lytic bacteriophages and a bacteriocin. Applied and Environmental Microbiology 69:4519-4526. 2003.

Materon, L.A. Survival of *Escherichia coli* O157:H7 applied to cantaloupes and the effectiveness of chlorinated water and lactic acid as disinfectants. World Journal of Microbiology and Biotechnology 19:867-873. 2003.

Saftner, R.A. et al. Sanitary dips with calcium propionate, calcium chloride, or a calcium amino acid chelate maintain quality and shelf stability of fresh-cut honeydew chunks. Postharvest Biology and Technology 29:257-269. 2003.

Sapers, G.M. and J.E. Sites. Efficacy of 1% hydrogen peroxide wash in decontaminating apples and cantaloupe melons. Journal of Food Science 68:1793-1797. 2003.

Suslow, T.V. et al. Production practices as risk factors in microbial food safety of fresh and fresh-cut produce. Comprehensive Reviews in Food Science and Food Safety 2(supplement):38-77. 2003.

Annous, B.A. et al. Surface pasteurization of whole fresh cantaloupes inoculated with *Salmonella* Poona or *Escherichia coli*. Journal of Food Protection 67:1876-1885. 2004.

Beuchat, L.R. and Scouten, A.J. Factors affecting survival, growth, and retrieval of *Salmonella* Poona on intact and wounded cantaloupe rind and in stem scar tissue. Food Microbiology 21:683-694. 2004.

Castillo, A. et al. *Salmonella* contamination during production of cantaloupe: A binational study. Journal of Food Protection 67:713-720. 2004.

Hammack, T.S. et al. Relative effectiveness of the Bacteriological Analytical Manual method for the recovery of *Salmonella* from whole cantaloupes and cantaloupe rinses with selected preenrichment media and rapid methods. Journal of Food Protection 67:870-877. 2004.

Karenlampi, R. and Hanninen, M.-L. Survival of *Campylobacter jejuni* on various fresh produce. International Journal of Food Microbiology 97:187-195. 2004.

Leverentz, B. et al. Optimizing concentration and timing of a phage spray application to reduce *Listeria monocytogenes* on honeydew melon tissue. Journal of Food Protection 67:1682-1686. 2004.

Li, Y. and Mustapha, A. Simultaneous detection of *Escherichia coli* O157:H7, *Salmonella*, and *Shigella* in apple cider and produce by a multiplex PCR. Journal of Food Protection 67:27-33. 2004.

Penteado, A.L. and Leitao, M.F.F. Growth of *Listeria monocytogenes* in melon, watermelon, and papaya pulps. International Journal of Food Microbiology 91:89-94. 2004.

Penteado, A.L. and Leitao, M.F.F. Growth of *Salmonella* Enteritidis in melon, watermelon and papaya pulp stored at different times and temperatures. Food Control 15:369-373. 2004.

Richards, G.M. and Beuchat, L.R. Attachment of *Salmonella* Poona to cantaloupe rind and stem scar tissues as affected by temperature of fruit and inoculum. Journal of Food Protection 67:1359-1364. 2004.

Richards, G.M. et al. Survey of yeasts for antagonistic activity against *Salmonella* Poona in cantaloupe juice and wounds in rinds coinfected with phytopathogenic molds. Journal of Food Protection 67:2132-2142. 2004.

Rodgers, S.L. et al. A comparison of different chemical sanitizers for inactivating *Escherichia coli* O157:H7 and *Listeria monocytogenes* in solution and on apples, lettuce, strawberries, and cantaloupe. Journal of Food Protection 67:721-731. 2004.

Sanglay, G.C. et al. Recovery of *Salmonella* spp. From raw produce surfaces using ultrasonication. Foodborne Pathogens and Disease 1:295-299. 2004.

Taitt, C.R. et al. Detection of *Salmonella enterica* serovar Typhimurium by using a raid, array-based immunosensor. Applied and Environmental Microbiology 70:152-158. 2004.

Ukuku, D.O. Effect of hydrogen peroxide treatment on microbial quality and appearance of whole and fresh-cut melons contaminated with *Salmonella* spp. International Journal of Food Microbiology 95:137-146. 2004.

Ukuku, D.O. and Fett, W.F. Effect of nisin in combination with EDTA, sodium lactate, and potassium sorbate for reducing *Salmonella* on whole and fresh-cut cantaloupe. Journal of Food Protection 67:2143-2150. 2004.

Ukuku, D.O. et al. Method of applying sanitizers and sample preparation affects recovery of native microflora and *Salmonella* on whole cantaloupe surfaces. Journal of Food Protection 67:999-1004. 2004.

Ukuku, D.O. et al. Effect of hot water and hydrogen peroxide treatments on survival of *Salmonella* and microbial quality of whole and fresh-cut cantaloupe. Journal of Food Protection 67:432-437. 2004.

Annous, B.A. et al. Improved recovery procedure for evaluation of sanitizer efficacy in disinfecting contaminated cantaloupes. Journal of Food Science 70:M242-M247. 2005.

Annous, B.A. et al. Biofilm formation by *Salmonella* spp. on cantaloupe melon. Journal of Food Safety 25:276-287. 2005.

Bastos, M.doS.R. et al. The effect of association of santizers and surfactant in the microbiota of the cantaloupe (*Cucumis melo* L.) melon surface. Food Control 16:369-373. 2005.

Duffy, E.A. et al. Concentrations of *Escherichia coli* and genetic diversity and antibiotic resistance profiling of *Salmonella* isolated from irrigation water, packaging shed equipment, and fresh produce in Texas. Journal of Food Protection 68:70-79. 2005.

Johnston, L.M. et al. A field study of the microbiological quality of fresh produce. Journal of Food Protection 69:1840-1847. 2005.

Kim, H. and Beuchat, L.R. Survival and growth of *Enterobacter sakazakii* on fresh-cut fruits and vegetables and in unpasteurized juices as affected by storage temperature. Journal of Food Protection 68:2541-2552. 2005.

Parnell, T.L. et al. Reducing *Salmonella* on cantaloupes and honeydew melons using practices applicable to postharvest handling, foodservice, and consumer preparation. International Journal of Food Microbiology 99:59-70. 2005.

Richards, G.M. and Beuchat, L.R. Infection of cantaloupe rind with *Cladosporium* caldosporioides and *Penicillium expansum*, and associated migration of *Salmonella* Poona in edible tissue. International Journal of Food Microbiology 103:1-10. 2005.

Richards, G.M. and Beuchat, L.R. Metabiotic associations of molds and *Salmonella* Poona on intact and wounded cantaloupe rind. International Journal of Food Microbiology 97:327-339. 2005.

Sharma, M. et al. Thermal tolerance of acid adapted and unadapted *Salmonella*, *Escherichia coli* O157:H7, and *Listeria monocytogenes* in cantaloupe juice and watermelon juice. Letters in Applied Microbiology 41:448-453. 2005.

Stine, S.W. et al. Effect of relative humidity on preharvest survival of bacterial and viral pathogens on the surface of cantaloupe, lettuce, and bell peppers. Journal of Food Protection 68:1352-1358. 2005.

Stine, S.W. et al. Application of microbial risk assessment to the development of standards for enteric pathogens in water used to irrigate fresh produce. Journal of Food Protection 68:913-918. 2005.

Ukuku, D.O. et al. Use of hydrogen peroxide in combination with nisin, sodium lactate and citric acid for reducing transfer of bacterial pathogens from whole melon surfaces to fresh-cut pieces. International Journal of Food Microbiology 104:225-233. 2005.

Ukuku, D.O. et al. ATP bioluminescence assay for estimation of microbial populations of fresh-cut melon. Journal of Food Protection 68:2427-2432. 2005.

Vargas, A.M. et al. Detection of fecal contamination on cantaloupes using hyperspectral fluorescence imagery. Journal of Food Science 70:E471-E476. 2005.

Bowen, A. et al. Infections associated with cantaloupe consumption: A public health concern. Epidemiology and Infection 134:675-685. 2006.

Boynton, B.B. et al. Effects of low-dose electron beam irradiation on respiration, microbiology, texture, color, and sensory characteristics of fresh-cut cantaloupe stored in modified atmosphere packages. Journal of Food Science 71:S149-S155. 2006.

Dipersio, P.A. et al. Sensory evaluation of home dried fruit prepared using treatments that enhance destruction of pathogenic bacteria. Journal of Food Quality 29:47-64. 2006.

Espinoza-Medina, I.E. et al. PCR identification of *Salmonella*: Potential contamination sources from production and postharvest handling of cantaloupes. Journal of Food Protection 69:1422-1425. 2006.

Fan, X. et al. Combination of hot-water surface pasteurization of whole fruit and low-dose gamma irradiation of fresh-cut cantaloupe. Journal of Food Protection 69:912-919. 2006.

Hammack, T.S. et al. Effect of sample preparation and preenrichment media on the recovery of *Salmonella* from cantaloupes, mangoes, and tomatoes. Journal of AOAC International 89:180-184. 2006.

Johnston, L.M. et al. A field study of the microbiological quality of fresh produce of domestic and Mexican origin. International Journal of Food Microbiology 112:83-95. 2006.

Solomon, E.B. et al. Thermal inactivation of *Salmonella* on cantaloupes using hot water. Journal of Food Science 71:M25-M30. 2006.

Ukuku, D.O. Effect of sanitizing treatments on removal of bacteria from cantaloupe surface, and recontamination with *Salmonella*. Food Microbiology 23:289-293. 2006.

Ukuku, D.O. and Fett, W.F. Effects of cell surface charge and hydrophobicity on attachment of 16 *Salmonella* serovars to cantaloupe rind and decontamination with sanitizers. Journal of Food Protection 69:1835-1843. 2006.

Ukuku, D.O. et al. Effect of vacuum-steam-vacuum treatment of microbial quality of whole and fresh-cut cantaloupe. Journal of Food Protection 69:1623-1629. 2006.

Wang, H. et al. Dual-phasic inactivation of *Escherichia coli* O157:H7 with peroxyacetic acid, acidic electrolyzed water and chlorine on cantaloupes and fresh-cut apples. Journal of Food Safety 26:335-347. 2006.

Wang, Z. et al. Influence of gamma irradiation of enzyme, microorganism, and flavor of cantaloupe (*Cucumis melo* L.) juice. Journal of Food Science 71:M215-M220. 2006.

Zhou, B. et al. 1-Methylcyclopropene counteracts ethylene-induced microbial growth on fresh-cut watermelon. Journal of Food Science 71:M180-M184. 2006.

Critzer, F.J. et al. Atmospheric plasma inactivation of foodborne pathogens on fresh produce surfaces. Journal of Food Protection 70:2290-2296. 2007.

Materon, L.A. et al. Identification of sources of microbial pathogens on cantaloupe rinds from pre-harvest to post-harvest operations. World Journal of Microbiology and Biotechnology 23:1281-1287. 2007.

Mosqueda-Melgar, J. et al. Influence of treatment time and pulse frequency on *Salmonella* ENTERITIDIS, *Eshcherichia coli* and *Listeria monocytogenes* populations inoculated in melon and watermelon juices treated by pulsed electric. International Journal of Food Microbiology 117:192-200. 2007.

Ukuku, D.O. and Sapers, G.M. Effect of time before storage and storage temperature on survival of *Salmonella* inoculated on fresh-cut melons. Food Microbiology 24:288-295. 2007.

Wang, H. et al. Modeling the effect of washing solution flow conditions on *Escherichia coli* O157:H7 population reduction on fruit surfaces. Journal of Food Protection 70:2533-2540. 2007.

Ailes, E.C. et al. Microbial concentrations on fresh produce are affected by postharvest processing, importation, and season. Journal of Food Protection 71:2389-2397. 2008.

Akins, E.D. et al. Washing practices on the microflora on Georgia-grown cantaloupes. Journal of Food Protection 71:46-51. 2008.

Bassett, J. and McClure, P. A risk assessment approach for fresh fruits. Journal of Applied Microbiology 104:925-943. 2008.

Belletti, N. et al. Antimicrobial efficacy of citron essential oil on spoilage and pathogenic microorganisms in fruit-based salads. Journal of Food Science 73:M331-M338. 2008.

Fan, X. et al. Effect of hot water surface pasteurization of whole fruit on shelf-life and quality of fresh-cut cantaloupe. Journal of Food Science 73:M91-M98. 2008.

Gallegos-Robles, M.A. et al. Identification of *Salmonella* serotypes isolated from cantaloupe and chile pepper production systems in Mexico by PCR–restriction fragment length polymorphism. Journal of Food Protection 71:2217–2222. 2008.

Mosqueda-Melgar, J. et al. Combination of high-intensity pulsed electric fields with natural antimicrobials to inactivate pathogenic microorganisms and extend the shelf-life of melon and watermelon juices. Food Microbiology 25:479-491. 2008.

Mahmoud, B.S.M. et al. Inactivation kinetics of inoculated *Escherichia coli* O157:H7, *Listeria monoctogenes*, and *Salmonella* Poona on whole cantaloupes by chlorine dioxide gas. Food Microbiology 25:857-865. 2008.

Munnoch, S.A. et al. A multi-state outbreak of *Salmonella* Saintpaul in Australia associated with cantaloupe consumption. Epidemiology and Infection 137:367-374. 2008.

Oms-Oliu, G. et al. Effect of superatmospheric and low oxygen modified atmospheres on shelf-life extension of fresh-cut melon. Food Control 19:191-199. 2008.

Perni, S. et al. Cold atmosphere plasma decontamination of the pericaps of fruit. Journal of Food Protection 71:302-308. 2008.

Perni, S. et al. Cold atmospheric plasma disinfection of cut fruit surfaces contaminated with migrating microorganisms. Journal of Food Protection 71:1619-1625. 2008.

Raybaudi-Massilia, R.M. et al. Edible alginate-based coating as carrier of antimicrobials to improve shelf-life and safety of fresh-cut melon. International Journal of Food Microbiology 121:313-327. 2008.

Selma, M.V. et al. Effect of gaseous ozone and hot water on microbial and sensory quality of cantaloupe and potential transference of *Escherichia coli* O157:H7 during cutting. Food Microbiology 25:162-168. 2008.

Selma, M.V. et al. Reduction by gaseous ozone of *Salmonella* and microbial flora associated with fresh-cut cantaloupe. Food Microbiology 25:558-565. 2008.

Silveira, A.C. et al. Alternative sanitizers to chlorine for use on fresh-cut "Galia" (*Cucumis melo* var. *catalupensis*) melon. Journal of Food Science 73:M405-M411. 2008.

Fan X. et al. Use of chemical sanitizers to reduce microbial populations and maintain quality of whole and fresh-cut cantaloupe. Journal of Food Protection 72:2453-2460. 2009.

Hanning, I.B. et al. Salmonellosis outbreaks in the United States due to fresh produce: Sources and potential intervention measures. Foodborne Pathogens and Disease 6:635-648. 2009.

Kim, J. et al. Evaluating the effect of environmental factors on pathogen regrowth in compost extract. Microbial Ecology 58:498-508. 2009.

Morales-Rayas, R. et al. Simultaneous separation and detection of hepatitis A virus and norovirus in produce. International Journal of Food Microbiology 139:48-55. 2010.

Munnoch, S. A. et al. A multi-state outbreak of *Salmonella* Saintpaul in Australia associated with cantaloupe consumption. Epidemiology and Infection 137:367-374. 2009.

Sharma, M. et al. Effectiveness of bacteriophages in reducing *Escherichia coli* O157:H7 on fresh-cut cantaloupes and lettuce. Journal of Food Protection 72:1481-1485. 2009.

Silagyi, K. et al. Production of biofilm and quorum sensing by *Escherichia coli* O157:H7 and its transfer from contact surfaces to meat, poultry, ready-to-eat deli, and produce products. Food Microbiology 26:514-519. 2009.

Wang, H. et al. Effect of surface roughness on retention and removal of *Escherichia coli* O157:H7 on surfaces of selected fruits. Journal of Food Science 74:E8-E15. 2009.

Issa-Zacharia, A. et al. A review of microbiological safety of fruits and vegetables and the introduction of electrolyzed water as an alternative to sodium hypochlorite solution. African Journal of Food Science 4:778-789. 2010.

Silveira, A.C. et al. Emerging sanitizers and clean room packaging for improving the microbial quality of fresh-cut 'Galia' melon. Food Control 21;863-871. 2010.

Anderson, M. et al. Pathogen-produce pair attribution risk ranking tool to prioritize fresh produce commodity and pathogen combinations for further evaluation (P³ARRT). Food Control 22:1865-1872. 2011.

Bolton, D.J. et al. Incidence and survival of non-O157 verocytotoxigenic *Escherichia coli* in soil. Journal of Applied Microbiology 111:484-490. 2011.

Chen, S. et al. Rapid detection of viable salmonellae in produce by coupling propidium monoazide with loop-mediated isothermal amplification. Applied and Environmental Microbiology 77:4008-4016. 2011.

Cosgrove et al. Multistate outbreak of listeriosis associated with Jensen Farms cantaloupe – United States, August – September 2011. Morbidity and Mortality Weekly Report 60:1-2. 2011.

Odjadjare, E.E.O. et al. Microbial and physicochemical quality of an urban reclaimed wastewater used for irrigation and aquaculture in South Africa. African Journal of Microbiology Research 5:2179-2186. 2011.

Abadias, M. et al. Growth potential of *Escherichia coli* O157:H7 on fresh-cut fruits (melon and pineapple) and vegetables (carrot and escarole) stored under different conditions. Food Control 27:37-44. 2012.

Chen, W. et al. Inactivation of *Salmonella* on whole cantaloupe by application of an antimicrobial coating containing chitosan and allyl isothiocyanate. International Journal of Food Microbiology 155:165-170. 2012.

Doyle, M.P. and M.C. Erickson. Opportunities for mitigating pathogen contamination during on-farm food production. International Journal of Food Microbiology 152:54-74. 2012.

Fishburn, J.D. et al. Efficacy of various consumer-friendly produce washing technologies in reducing pathogens on fresh produce. Food Protection Trends 32:456-466. 2012.

Harris, L.J. et al. A framework for developing research protocols for evaluation of microbial hazards and controls during production that pertain to the quality of agricultural water contacting fresh produce that may be consumed raw. Journal of Food Protection 75:2251-2273. 2012.

Hoelzer, K. et al. *Listeria monocytogenes* growth dynamics on produce: A review of the available data for predictive microbiology. Foodborne Pathogens and Disease 9:661-673. 2012.

Lewis Ivey, M.L. et al. Vegetable producers' perceptions of food safety hazards in the Midwestern USA. Food Control 26:453-465. 2012.

Mahmoud, B.S.M. Effects of X-ray treatments on pathogenic bacteria, inherent microflora, color, and firmness on whole cantaloupe. International Journal of Food Microbiology 156:296-300. 2012.

Moreira, R.G., A.F. Puerta-Gomez, J. Kim, and M.E. Castell-Perez. Factors affecting radiation D-values (D10) of an *Escherichia coli* cocktail and *Salmonella* Typhimurium LT2 inoculated in fresh produce. Journal of Food Science 71:E104-E111.

Moreira, R.G. et al. Factors affecting radiation D-values (D10) of an *Escherichia coli* cocktail and *Salmonella* Typhimurium LT2 inoculated in fresh produce. Journal of Food Science 71:E104-E111.

Olaimat, A.N. and R.A. Holley. Factors influencing the microbial safety of fresh produce: A review. Food Microbiology 32:1-19. 2012.

Park, S. et al. Risk factors for microbial contamination in fruits and vegetables at the preharvest level: A systematic review. Journal of Food Protection 75:2055-2081. 2012.

Parker, J.S. et al. An expert guide to understanding grower decisions related to fresh fruit and vegetable contamination prevention and control. Food Control 26:107-116. 2012.

Soon, J.M. and R.N. Baines. Food safety training and evaluation of handwashing intention among fresh produce farm workers. Food Control 23:437-448. 2012.

Ukuku, D.O. et al. Effect of native microflora, waiting period, and storage temperature on *Listeria monocytogenes* serovars transferred from cantaloupe rind to fresh-cut pieces during preparation. Journal of Food Protection 75:1912-1919. 2012.

Vadlamudi, S. et al. Effect of chemical sanitizers on *Salmonella enterica* serovar Poona on the surface of cantaloupe and pathogen contamination of internal tissues as a function of cutting procedure. Journal of Food Protection 75:1766-1773. 2012.

Verrill, L. et al. Consumer vegetable and fruit washing practices in the United States, 2006 and 2010. Food Protection Trends 32:164-172. 2012.

Annous, B.A. et al. Commercial thermal process for inactivating *Salmonella* Poona on surfaces of whole fresh cantaloupes. Journal of Food Protection 76:420-428. 2013.

Bae, D. et al. Characterization of a potential *Listeria monocytogenes* virulence factor associated with attachment to fresh produce. Applied and Environmental Microbiology 79:6855-6861. 2013,

Bihn, E.A., et al. Use of surface water in the production of fresh fruits and vegetables: A survey of fresh produce growers and their water management practices. Food Protection Trends 33:307-314. 2013.

Feng, P.C.H. and Reddy, S. Prevalences of Shiga toxin subtypes and selected other virulence factors among Shiga-toxigenic *Escherichia coli* strains isolated from fresh produce. Applied and Environmental Microbiology 79:6917-6923. 2013.

Gombas, D.E. Produce GAPs harmonization: The goal is in sight. Food Safety Magazine 19(3):58, 60-64, & 66. 2013.

Li, D. et al. Development and validation of a mathematical model for growth of pathogens in cut melons. Journal of Food Protection 76:953-958. 2013.

Massey, L.M. et al. Electrostatic spray of food-grade organic acids and plant extract to reduce *Escherichia coli* O157:H7 on fresh-cut cantaloupe cubes. Journal of Food Safety 33:71-78. 2013.

McCollum, J.T. et al. Multistate outbreak of listeriosis associated with cantaloupe. New England Journal of Medicine 369:944-953. 2013

Park, M.-K. et al. The analytical comparison of phage-based magnetoelastic biosensor with TaqMan-based quantitative PCR method to detect *Salmonella* Typhimurium on cantaloupes. Food Control 33:330-336. 2013.

Sospedra, I. et al. Survey of microbial quality of plant-based foods served in restaurants. Food Control 30:418-422. 2013.

Strawn, L.K. et al. Landscape and meteorological factors affecting prevalence of three food-borne pathogens in fruit and vegetable farms. Applied and Environmental Microbiology 79:588-600. 2013.

Trinetta, V. et al. The application of high-concentration short-time chlorine dioxide treatment for selected speciality crops including Roma tomatoes (*Lycopersicon esculentum*), cantaloupes (*Cucumis melo* ssp. *melo* var. *cantaloupensis*) and strawberries (*Fragaria x ananassa*). Food Microbiology 34:296-302. 2013.

Wang, Q. et al. Physical removal and transfer of murine norovirus and hepatitis A virus from contaminated produce by scrubbing and peeling. Journal of Food Protection 76:85-92. 2013.

Webb, C.C. et al. Evaluation of levulinic acid and sodium dodecyl sulfate as a sanitizer for use in processing Georgia-grown cantaloupes. Journal of Food Protection 76:1767-1772. 2013.

Yao, Z. et al. Survival of *Escherichia coli* O157:H7 in soils from vegetable fields with different cultivation patterns. Applied and Environmental Microbiology 79:1755-1756. 2013.

Byrne, L. et al. A multi-country outbreak of *Salmonella* Newport gastroenteritis in Europe associated with watermelon from Brazil, confirmed by whole genome sequencing: October 2011 to January 2012. Eurosurveillance 19(31):07 August (online). 2014.

Danyluk, M.D. et al. Modeling the growth of *Listeria monocytogenes* on cut cantaloupe, honeydew and watermelon. Food Microbiology 38:52-55. 2014.

EFSA. Scientific Opinion on the risk posed by pathogens in food of non-animal origin. Part 2 (Salmonella in melons). EFSA Journal 12(10):3831 (Online). 2014.

Erickson, M.C. et al. Thermal and nonthermal factors affecting survival of *Salmonella* and *Listeria monocytogenes* in animal manure-based compost mixtures. Journal of Food Protection 77:1512-1518. 2014.

Feng, P.C.H. and S.P. Reddy. Prevalence and diversity of enterotoxigenic *Escherichia coli* strains in fresh produce. Journal of Food Protection 77:820-823. 2014.

Hoelzer, K. et al. Reduction of *Listeria monocytogenes* contamination on produce – A quantitative analysis of common liquid fresh produce wash compounds. Food Control 46:430-440. 2014.

Hong, Y.-K. et al. Predictive modeling for growth of non- and cold-adapted *Listeria monocytogenes* on fresh-cut cantaloupe at different storage temperatures. Journal of Food Science 79:M1168-M1174. 2014.

Jones, L.A. et al. Plant-pathogenic oomycetes, *Escherichia coli* strains, and *Salmonella* spp. frequently found in surface water used for irrigation of fruit and vegetables crops in New York State. Applied and Environmental Microbiology 80:4814-4820. 2014.

Martinon, M.E. et al. Development of a multilayered antimicrobial edible coating for shelf-life extension of fresh-cut cantaloupe (*Cucumis melo* L.) stored at 4°C. LWT – Food Science and Technology 56:341-350. 2014.

Melotto, M. et al. Plant innate immunity against human bacterial pathogens. Frontiers in Microbiology 5:article 411 (online). 2014.

Russo, P. et al. A fast, reliable, and sensitive method for detection and quantification of *Listeria monocytogenes* and *Escherichia coli* O157:H7 in ready-to-eat fresh-cut products by MPN-qPCR. BioMed Research International 2014:608296 (Online). 2014.

Upadhyay, A. et al. Efficacy of plant-derived compounds combined with hydrogen peroxide as antimicrobial wash and coating treatment for reducing *Listeria monocytogenes* on cantaloupes. Food Microbiology 44:47-53. 2014.

van Overbeek, L.S. et al. The arable ecosystem as battleground for emergence of new human pathogens. Frontiers in Microbiology 5:article 104 (online). 2014.

Walsh, K.A., S.D. Bennett, M. Mahovic, and L.H. Gould. Outbreaks associated with cantaloupe, watermelon, and honeydew in the United States, 1973-2011. Foodborne Pathogens and Disease 11:945-952. 2014.

Yeni, F. et al. Alpas. Rapid and standardized methods for detection of foodborne pathogens and mycotoxins on fresh produce. Food Control 40:359-367. 2014.

Callejon, R.M. et al. Reported foodborne outbreaks due to fresh produce in the United States and European Union: Trends and causes. Foodborne Pathogens and Disease 12:32-38. 2015.

Dundore-Arias, J.P. et al. Influence of *prgH* on the persistence of ingested *Salmonella enterica* in the leafhopper *Macrosteles quadrilineatus*. Applied and Environmental Microbiology 81:6345-6354. 2015.

Erickson, M.C. et al. Contamination of knives and graters by bacterial foodborne pathogens during slicing and grating of produce. Food Microbiology 52:138-145. 2015.

Giangaspero, A. et al., Molecular detection of *Cyclospora* in water, soil, vegetables and humans in southern Italy signals a need for improved monitoring by health authorities. International Journal of Food Microbiology 211:95-100. 2015.

Heredia, N. et al. Validation of a novel rinse and filtration method for efficient processing of fresh produce samples for microbiological indicator enumeration. Journal of Food Protection 78:525-530. 2015.

Hou, M.A. et al. Food safety standards and international supply chain organization: A case study of the Moroccan fruit and vegetable exports. Food Control 55:190-199. 2015.

Huang, J. et al. Growth of *Salmonella enterica* and *Listeria monocytogenes* on fresh-cut cantaloupe under different temperature abuse scenarios. Journal of Food Protection 78:1125-1131. 2015.

Laury-Shaw, A. et al. Current trends in food safety practices for small-scale growers in the Midwest. Food Protection Trends 35:461-469. 2015.

Liu, N.T. et al. Effects of environmental parameters on the dual-species biofilms formed by *Escherichia coli* O157:H7 and *Ralstonia insidiosa*, a strong biofilm producer isolated from a fresh-cut produce processing plant. Journal of Food Protection 78:121-127. 2015.

Nuesch-Inderbinen, M. et al. Assessment of the prevalence of extended-spectrum Beta-lactamase-producing *Enterobacteriaceae* in ready-to-eat salads, fresh-cut fruit, and sprouts from the Swiss market. Journal of Food Protection 78:1178-1181. 2015.

Palekar, M.P. et al. Reduction of *Salmonella enterica* serotype Poona and background microbiota on fresh-cut cantaloupe by electron beam irradiation. International Journal of Food Microbiology 202:66-72. 2015.

Ramos-Villarroel, A.Y. et al. Combined effects of malic acid dip and pulsed light treatments on the inactivation of *Listeria innocua* and *Escherichia coli* on fresh-cut produce. Food Control 52:112-118. 2015.

Raede, J. *Listeria* guidance & best practices in produce facilities. Food Safety Magazine 21(1):58-63. 2015.

Ukuku, D.O. et al. Efficacy of sanitizer treatments on survival and growth parameters of *Escherichia coli* O157:H7, *Salmonella*, and *Listeria monocytogenes* on fresh-cut pieces of cantaloupe during storage. Journal of Food Protection 78:1288-1295. 2015

Weller, D. et al. Irrigation is significantly associated with an increased prevalence of *Listeria monocytogenes* in produce production environments in New York State. Journal of Food Protection 78:1132-1141. 2015.

Wyard, G. and N. Lewis. A long row to hoe for safer food. Food Safety Magazine 21(1):50-57. 2015.

Yoo, B.K. et al. Growth characteristics of Shiga toxin-producing *Escherichia coli* (STEC) stressed by chlorine, sodium chloride, acid, and starvation on lettuce and cantaloupe. Food Control 55:97-102. 2015.

Zaczek, M., et al. Phages in the global fruit and vegetable industry. Journal of Applied Microbiology 118:537-556. 2015.

Zhang, Y. et al. Effect of algenate coatings with cinnamon bark oil and soybean oil on quality and microbiological safety of cantaloupe. International Journal of Food Microbiology 215:25-30. 2015.

Martiniez, M.R. et al. Capacity of Listeria monocytogenes strains from the 2011 cantaloupe outbreak to adhere, survive, and grow on cantaloupe. Journal of Food Protection 79:757-763. 2016.

Garner, D. and S. Kathariou. Fresh produce-associated listeriosis outbreaks, sources of concern, teachable moments, and insights. Journal of Food Protection 79:337-344. 2016.

Hadjilouka, A. et al. Expression of *Listeria monocytogenes* key virulence genes during growth in liquid medium, on rocket and melon at 4, 10, and 30 C. Food Microbiology 55:7-15. 2016.

Lone, A. et al. Development of prototypes of bioactive packaging materials based on immobilized bacteriophages for control of growth of bacterial pathogens in foods. International Journal of Food Microbiology 217:49-58. 2016.

Lopez, K. et al. Investigation of handling practices for fresh produce and the efficacy of commercially available produce washes on removal of pathogens and natural microflora on whole cantaloupe surfaces. Food Control 68:251-259. 2016

Santo, D. et al. Survival and growth of *Cronobacter sakazakii* on fresh-cut fruit and the effect of UV-C illumination and electrolyzed water in the reduction of its population. International Journal of Food Microbiology 231:10-15. 2016.

Shearer, A.E.H. et al. Transfer of pathogens from cantaloupe rind to preparation surfaces and edible tissue as a function of cutting method. Journal of Food Protection 79:764-770. 2016.

Spadafora, N.D. et al. Detection of *Listeria monocytogenes* in cut melon fruit using analysis of volatile organic compounds. Food Microbiology 54:52-59. 2016.

Svoboda, A. et al. Effectiveness of broad-spectrum chemical produce sanitizers against foodborne pathogens as in vitro planktonic cells and on the surface of whole cantaloupes and watermelons. Journal of Food Protection 79:524-530. 2016.

Upadhyay, A. et al. Inactivation of *Listeria monocytogenes*, *Salmonella* spp. and *Escherichia coli* O157:H7 on cantaloupes by octenidine dihydrochloride. Food Microbiology 58:121-127. 2016.

Willis, C. et al. Assessment of the microbiological safety of precut fruit from retail and catering premises in the United Kingdom. Journal of Food Protection 79:598-604. 2016.